

TURF MAX MSMA 6+

SELECTIVE POSTEMERGENCE GRASS HERBICIDE FOR TURF AND NON-CROPLAND AREAS

Prokoz Turf Max MSMA 6+ is a postemergence grass herbicide used to control weeds, particularly grassy weeds. It is used to control Bahiagrass, Barnyardgrass, chickweed, cocklebur, crabgrass (large & smooth), foxtail, Goosegrass, Johnsongrass, pigweed, nutsedge, punturevine, sandbur, and watergrass to name a few. Some of the sites Prokoz Turf Max 6+ can be applied to include turf, golf courses, grass seed crops, forestry, and non-crop areas. A surfactant has already been added so there is no need to mix a wetting agent to your spray tank, saving you time and money. You will need to apply during warm weather, 80 to 90 degrees, and when weeds are in an active stage of growth to obtain optimum results.

Active Ingredient

MSMA (Monosodium Acid Methanearsonate) 47.8% / 6 lbs. per gallon active

Packaging

4 x 1 GL / CS
2 x 2.5 GL / CS

Application Sites

Turfgrass, ornamental turfgrass, grass seed crops, non-bearing fruit and nuts, citrus (bearing and non-bearing), forestry, and non-cropland areas including, drainage ditch banks, right-of-ways, fence rows, and storage yards

Application Rates

Turf – 2 to 2 2/3 pints in 40 to 100 gallons of water / acre with thorough coverage.
Spot treatment rate – 1 to 2 oz. in 5 gallon water / 1000 sq. ft. depending on turf type.
Non-turf rates vary, please refer to label.

Benefits

- Postemergent control of tough grassy weeds such as foxtail, sandbur, and large and smooth crabgrass.
- Turf can be reseeded two weeks after application.
- Can be used in the control of labeled conifers.
- Surfactant already added. Saves time and money.
- Completely water soluble.

Important

For turf apply when temperatures are between 80 and 90 degrees F. Do not water for at least 24 hours. Have turf cut to 1 1/2 inches and still actively growing. Bermudagrass, Bluegrass, and Zoysiagrass have shown tolerance to MSMA applications, still some temporary discoloring may occur. Injury may occur if applied to Fescues and Bentgrasses. Do not apply to St. Augustinegrass, Carpetgrass, Centipedegrass, or Dichondra lawns.